WHAT ARE THE WARNING SIGNS OF BLOOD CLOTS IN THE LEG?



- 1- Swelling of the leg or calf muscle
- 2- Pain that may worsen when standing or walking
- 3- Warmth and redness of the leg

WHAT ARE THE WARNING SIGNS OF BLOOD CLOTS IN THE LUNG?

- 1- Unexplained sudden breathlessness
- 2- Sudden sharp chest pain, often when breathing or straining
- 3- Coughing, or coughing up blood
- 4- Feeling a very fast heart rate, even at rest



HOW TO PREVENT A BLOOD CLOT IN PREGNANCY AND AFTER CHILDBIRTH?



Light exercise -Walk for 30 mins per day



Avoid alcohol



Stop smoking



Drink 6-8 cups of water a day



Healthy diet



If planning to travel ≥ 4 hrs please seek medical advice.

Follow us f 🔰 🌀 🖺 www.phcc.qa



PREVENTING BLOOD CLOTS

In pregnancy and after childbirth



DEFINITION:


Venous Thromboembolism (VTE) is the medical term for a condition in which a blood clot starts in a vein in the body.

TYPES OF BLOOD CLOT:

If this occurs in the leg, it is called a **DVT** (deep vein thrombosis).



If it occurs in the lung, it is called a **PE** (pulmonary embolism).



KNOW YOUR RISK:

Pregnancy and childbirth can increase the risk for blood clots

A woman is at increased risk up to 6 weeks after her baby is born

Risks for blood clots during pregnancy, childbirth and up to 6 weeks after the baby is born include:



Prolonged immobility due to bed rest during pregnancy or recovery after delivery



A previous blood clot, a family history of clots, or a genetic blood clotting disorder



Delivery by cesarean section



Pregnancy at age 35 or older



Overweight or Obesity BMI≥30



Smoking



Varicose Veins



Certain pregnancy complications, such as pregnancy with multiple babies or the use of fertility treatments involving hormones

If you have any of these risk factors, or are unsure, please speak to your doctor urgently.